

OBSERVING THE UNOBSERVABLE: CORONAL LINE DATA FROM NARROW-LINE SEYFERT 1S AS A TEST OF EUV CONTINUUM MODELS

P.J. Bleackley and P.T.O'Brien

Department of Physics and Astronomy, University of Leicester

ABSTRACT We have measured coronal line emission for a sample of 19 Narrow-Line Seyfert 1 galaxies. Kinematic analysis shows that the Coronal Line Region (CLR) consists of gas in a decelerating outflow.

We intend to use these data as a test of the EUV emission of NLS1s. By comparison with photoionization modelling for a number of power-law + black body spectra, we plan to deduce the most likely EUV continua for these objects as well as explore physical conditions in the CLR.

KEYWORDS: galaxies active — galaxies Seyfert — spectroscopy coronal lines — AGN outflows — AGN soft excess

1. INTRODUCTION

Narrow-line Seyfert 1 objects (NLS1s) are a phenomenologically interesting subclass of Seyfert 1s. As their name suggests, they have the narrowest permitted lines of all Seyfert 1s ($\text{FWHM}(\text{H}\beta) \lesssim 2000 \text{ km s}^{-1}$), which appear to be associated with a range of other extremal properties, including strong, steep, soft X-ray excesses, strong permitted Fe II lines, weak forbidden lines, and a high degree of short timescale variability in X-rays. Coronal lines, forbidden transitions from highly ionised states of iron, are often present in their optical spectra. Their properties form a continuum with those of “normal” Seyfert 1s, but as they show extremal properties, they are important for testing AGN unification schemes. It is thought likely that they are Seyfert 1s with low mass black holes and/or high accretion rates.

To further understand the nature of the soft UV/X-ray excess, which is an important component of the energy output of these objects, we wish to study the EUV continuum. As this is not directly observable for extragalactic objects due to Galactic absorption, we use coronal lines (which have ionisation energies $\sim$ a few hundred eV) as diagnostics of this region. This approach also allows us to investigate the physical conditions in the coronal line region, particularly by kinematic methods.

2. OBSERVATIONS AND DATA REDUCTION

Nineteen NLS1s from the soft X-ray observed sample of Boller et al. (1996) were observed between 5 – 11 August 1996, using the Intermediate Dispersion Spectrograph mounted on the Isaac Newton Telescope at La Palma, obtaining dispersions of $1.6 - 3.2 \text{ \AA pixel}^{-1}$. The data were reduced with the Starlink project’s **FIGARO** software, and fitted using the **ELF** routines in **DIPS0**, which simultaneously fitted gaussian line profiles and a linear continuum. This approach allowed the deblending of neighbouring lines.

The coronal lines [Fe VII] $\lambda 6038.9$, [Fe X] $\lambda 6374.6$, [Fe XI] $\lambda 7891.9$ and [Fe XIV] $\lambda 5302.86$ were identified where possible, and upper limits were placed on them otherwise. Line profiles were then transformed into velocity space relative to the rest-frame of the [O III] $\lambda\lambda 4959, 5007$ doublet.

3. RESULTS




FIGURE 1. The decelerating outflow of the Coronal Line Region

3.1. Kinematic analysis

Erkens et al. (1997) noted a tendency for broader coronal lines to be more blueshifted than narrower ones. To further investigate this correlation in a manner independent of black hole mass, we assume that the FWHM of coronal lines is governed by the virial theorem and define a *radial estimator*

$$\mathcal{R} = \frac{r_{[\text{Fe}]}}{r_{[\text{O III}]}} = \left(\frac{\text{FWHM}_{[\text{O III}]}}{\text{FWHM}_{[\text{Fe}]}} \right)^2$$

where $r_{[\text{Fe}]}$ is the distance from the central black hole at which a line, [Fe], is emitted. Plotting the outflow velocity δv against $\mathcal{R}$ shows a decelerating outflow, which was fitted to first order by $\delta v = (806 \pm 93)\mathcal{R} - (417 \pm 25) \text{ km s}^{-1}$ (see Figure 1). The minimum value of $\mathcal{R}$ observed is 0.024, which constrains the maximum size of the region over which the gas is accelerated.

For those objects for which at least 3 coronal lines had been detected, we plotted $\mathcal{R}$ against the ionisation potential, in order to determine if stratification of the CLR was important. For Arakelian 564, there is a trend towards more highly ionised gas being detected closer to the nucleus, but this trend is not unambiguously detected in other objects (see Figure 2).




FIGURE 2. Radial estimator against ionisation potential for coronal lines.

3.2. Comparison with X-ray data

Equivalent widths of [Fe VII] and [Fe X] were compared with ROSAT PSPC photon indices obtained from Boller et al. (1996) (see Figure 3). No strong correlation is present for either line.

4. CONCLUSIONS

Gas in the coronal line region is outflowing, and this outflow appears to be decelerating. The coronal line gas has reached an outflow velocity of 420 km s^{-1} within a distance from the nucleus estimated to be $< 1/40$ that of the NLR. Radiation pressure does not appear to drive motion in the CLR, as this would be expected to produce accelerating, rather than decelerating outflows (see Binette (1998) for an example of a radiation pressure driven model).




FIGURE 3. Equivalent width of [Fe VII] (crosses) and [Fe X] (triangles) against ROSAT PSPC photon index.

Coronal line strengths do not follow a simple correlation with ROSAT PSPC photon index. To investigate the EUV continuum in more detail, we plan to create photoionisation models based on a fixed power-law, with a black body component of variable luminosity and temperature added, and test the predictions of these models against our observed coronal line strengths. The models will also allow for stratification of the CLR.

ACKNOWLEDGEMENTS

We thank Dr. T. Roberts for his help with the observations. This work uses data obtained using the Isaac Newton Telescope, Observatorio de la Roque de los Muchachos, La Palma. P.J. Bleackley has been supported by a PPARC research studentship.

REFERENCES

- Binette, L., 1998, MNRAS, 294, L47–L51
- Boller, Th., Brandt, W.N., and Fink, H., 1996, A&A, 305, 53–73
- Erkens, U., Appenzeller, I., and Wagner, S., 1997, A&A, 323, 707–716